Graphs and Images

From Research to Quality Improvement: Adolescent NG/CT Screening and Lessons Learned from a Texas Academic Medical Center

Amanda Brosnan, MPAS, PA-C, Dr. Maria Trent, MD, MPH, FAAP, FSAHM, Dr. Stacey Griner, PhD, MPH, CPH, RDH




Figure 1: Methods and Results Overview

 Table 1. EMR Variables Requested for Analysis.

General Parameters	Date of visit
	Notation if a visit was a well-
	check
	Patient height and weight, date
	of birth, and age
	Clinic Location and medical
	provider seen
	Payer source
	Patient race, ethnicity, sex,
	language and zip code
Sexual Health Parameters	NG*, CT*, syphilis and HIV
	screening
	NG diagnosis*
	CT diagnosis*
	HIV diagnosis
	HPV vaccine received
	Pregnancy testing
	Contraceptive Counseling
	Sexually active? (Y/N)*
	Practicing safe sex?*
	(Y/N/Sometimes)

^{*}Indicates parameters of interest throughout research studies

Table 2. Data Results from Study 1 and Data Refinement for Study 2.

Study 1: Preliminary Needs Assessment		
Data Results	.1% proportion of patient visits screened for GC/CT	
Data Refinement (Moving to Study 2)	Reassess ways to capture data apart from ICD-10 codes	

 Table 3. Data Results from Study 2 and Data Refinement for Study 3.

Study 2: Data Investigation		
Data Results	3% proportion of patient visits screened for GC/CT	
	<0.2% patient visits with diagnoses of GC/CT	
	6% of patient visits with sexual activity documentation	
	3% of patient visits with safe sex practices documentation	
Data Refinement	Resolving discrepancies in best ways to capture data	
(Moving to Study 3)	Plan a series of informal unstructured interviews to improve understanding of data and system processes	




Figure 2. Discrepancies in Specimen Processing by Payor Source




Figure 3: Adolescent Sexual Health Data Collection and Reporting: Program Improvement